

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

Sh 09/320,323



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 41/10		A1	(11) International Publication Number: WO 96/13163
			(43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/GB95/02461 (22) International Filing Date: 18 October 1995 (18.10.95) (30) Priority Data: 08/333,442 1 November 1994 (01.11.94) US (71) Applicant: ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors: ENSMINGER, Michael, Paul; 22 Canterbury Court, Petaluma, CA 94954 (US). SHRIBBS, John, Martin; 108 Wyndham Way, Petaluma, CA 94954 (US). (74) Agents: RICKS, Michael, James et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG12 6YA (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>	

(54) Title: SELECTIVE CORN HERBICIDE

(57) Abstract

A method of selectively controlling undesirable vegetation in corn by applying an herbicidally effective amount of 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione to the locus of such vegetation.

=US 5,506,195

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

SELECTIVE CORN HERBICIDE

Field of the Invention

The present invention is directed to the use of 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione as a selective preemergence and postemergence herbicide in corn.


Background of the Invention

The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem researchers in the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use.

Unfortunately, many of such herbicides will exhibit phytotoxicity to the desired crop as well as to the weeds sought to be controlled. Thus, there is a long-standing need for selective herbicides which will control frequently occurring weeds but which will not adversely affect the crop plants when applied at herbicidally effective levels.

United States Patent 5,006,158 to Carter et al. discloses 2'-nitro-substituted benzoyl cyclohexanediones having the structure:

-2-



wherein R^1 and R^3 - R^6 are hydrogen or alkyl, R^2 is hydrogen, alkyl or alkoxy-carbonyl; R^7 is hydrogen or alkoxy; and R^8 is hydrogen, halogen, alkoxy, alkyl, OCF_3 , cyano, nitro, halo-alkyl, optionally substituted amino, optionally substituted aminosulfonyl, alkylcarbonyl, alkoxy-carbonyl or $R^9(SO)_n$ wherein n is 0, 1 or 2 and R^9 is substituted alkyl, phenyl or benzyl. Specifically disclosed, as Compound 26D, is 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione. Such compound is demonstrated to be an effective preemergence and postemergence herbicide against a wide variety of grasses, broadleaf weeds and sedges when applied at a rate of 4.48 kg/ha.

It has now been discovered that such compound will effectively control a broad range of weeds typically associated with corn without exhibiting any substantial phytotoxic effect on the corn itself.

Description of the Invention

This invention is directed to a method of selectively controlling undesirable vegetation in corn, comprising application of an herbicidally effective amount of 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione (hereinafter NMSC) to the locus of such vegetation. As is employed herein, the term "herbicide" is used to denote a compound which controls or modifies the growth of plants. The term "herbicidally effective amount" is used to indicate the quantity of such compound which is capable of producing a

controlling or modifying effect. Controlling or modifying effects include all deviations from natural development, for example: killing, retardation, leaf burn, albinism, dwarfing, and the like. The term "plants" refers to all physical parts of a plant, including seeds, seedlings, saplings, roots, tubers, stems, stalks, foliage, and fruits.

NMSC is a known compound and may be produced by methods such as those described in United States Patent 5,006,158 to Carter et al.

In the practice of the present invention, NMSC is applied to the locus of the vegetation to be controlled. Application rates will depend on the particular plant species and degree of control desired. In general, application rates of between about 1 and about 1,000 g/ha may be employed, with rates of between about 20 and about 500 g/ha being preferred.

NMSC can be formulated in the same manner in which herbicides are generally formulated. The object of the formulation is to apply the compositions to the locus where control is desired by a convenient method. The "locus" is intended to include soil, as well as established vegetation.

In practice, NMSC is applied as a formulation containing various adjuvants and carriers known to or used in the industry for facilitating dispersion. The choice of formulation and mode of application for NMSC may affect its activity, and selection will be made accordingly. The NMSC may thus be formulated as wettable powders, as flowable formulations, as emulsions, as granular formulations, as water dispersible granules, as powders or dusts, or as controlled release forms such as microcapsules. These formulations may contain as little as about 0.5% to as much as about 95% or more by weight of NMSC. The optimum amount in any particular

-4-

formulation will depend upon the materials in the formulation and the type of seeds or plants to be controlled.

Wettable powders (WP) are in the form of finely divided particles which disperse readily in water or other liquid carriers. The particles contain the active ingredient retained in a solid matrix. Typical solid matrices include fuller's earth, kaolin clays, silicas, and other readily wet organic or inorganic solids. Wettable powders normally contain about 5% to about 95% of the active ingredient plus a small amount of wetting, dispersing, or emulsifying agent.

Flowable formulations, also known as Suspension Concentrates (SC), are stable suspensions of active ingredient(s) intended for dissolution before use. This type of formulation consists of a dispersion of finely divided active ingredient(s) which may be practically insoluble or only slightly soluble in water or in organic solvents chosen as the diluent. It also contains inert materials such as dispersants, wetting agents, suspending aids and diluent. In general, flowable formulations tend to be creamy in appearance and readily mixable with water.

Emulsions, also known as emulsifiable concentrates (EC), are heterogeneous dispersions of one liquid in another liquid with which it is incompletely miscible. There are two common types, a dispersion of fine globules of an organic liquid in water (O/W type), and, less commonly, a dispersion of globules of an aqueous liquid in oil (W/O type). A stable mixture is produced by the addition of appropriate emulsifying agents. Typically, this type of formulation will contain 1 to 90% active ingredient.

Granular formulations, generally referred to as Impregnated Granular Formulations (GR) contain active ingredients impregnated in carriers. Inert materials of

-5-

granular formulations include extrudates, relatively coarse particles ("carriers"), and surface active agents. Typical carriers of granular formulations include: sand, fuller's earth, vermiculite, perlite, and other organic or inorganic materials which can be coated with the active compound. Typical surface active agents are: 1) heavy aromatic naphthas, kerosene and other petroleum fractions, 2) vegetable oils, and 3) stickers, such as dextrans, glue, or synthetic resins. Granular formulations are usually applied to weeds without being diluted.

Water Dispersible Granules (WG or WDG) formulations consist of small granules to be disintegrated and dispersed in water prior to application. Granules can be formed either by agglomeration or through the use of elevated pressure, for example, extrusion. Surface active agents, such as dispersants and wetting agents, are essential ingredients of the formulation. Clays, silicas and starch, among others, can be used as carriers.

Powders or dusts are free-flowing admixtures of the active ingredient with finely divided solids such as talc, clays, flours, and other organic and inorganic solids which act as dispersants and carriers.

Microcapsules are typically droplets or granules of the active material enclosed in an inert porous shell which allows escape of the enclosed material to the surroundings at controlled rates. Encapsulated droplets are typically about 1 to 50 microns in diameter. The enclosed liquid may include solvent in addition to the active compound. Encapsulated granules are generally porous granules with porous membranes sealing the granule pore openings, retaining the active species in liquid form inside the granule pores. Granules typically range from 1 millimeter to 1 centimeter, preferably 1 to 2 millimeters in diameter. Granules are formed by

-6-

extrusion, agglomeration or prilling, or are naturally occurring. Examples of such materials are vermiculite, sintered clay, kaolin, attapulgite clay, sawdust, and granular carbon. Shell or membrane materials include natural and synthetic rubbers, cellulosic materials, styrene-butadiene copolymers, polyacrylonitriles, polyacrylates, polyesters, polyamides, polyureas, polyurethanes, and starch xanthates.

Other useful formulations for herbicidal applications include simple solutions, sometimes described as flowable formulations, of the active ingredient in a solvent in which it is completely soluble at the desired concentration, such as water, acetone, alkylated naphthalenes, xylene, and other organic solvents. Pressurized sprayers, wherein the active ingredient is dispersed in finely-divided form as a result of vaporization of a low boiling dispersant solvent carrier, may also be used.

Many of these formulations include wetting, dispersing or emulsifying agents. Examples of these agents are alkyl and alkylaryl sulfonates and sulfates and their salts, polyhydric alcohols, polyethoxylated alcohols, esters, and fatty amines. These agents, when used, normally comprise from 0.1% to 15% by weight of the formulation.

Each of the above formulations can be prepared as a package containing the herbicide together with other ingredients of the formulation (diluent, emulsifiers, surfactants etc.). The formulations can also be prepared by a tank mix method in which the ingredients are obtained separately and combined at the grower site.

These formulations can be applied by conventional methods to the areas where control is desired. Dust and liquid compositions, for example, can be applied by the use of power dusters, boom and hand sprayers, and spray dusters. The

-7-

formulations can also be applied from airplanes as a dust or a spray or by rope wick applications. To modify or control growth of germinating seeds or emerging seedlings, dust and liquid formulations can be distributed in the soil to a depth of at least one centimeter below the soil surface or can be applied to the soil surface only, by spraying or sprinkling. The formulations can also be applied by addition to irrigation water, permitting penetration of the formulations into the soil together with the irrigation water. Dust compositions, granular compositions, or liquid formulations applied to the surface of the soil can be distributed below the surface of the soil by conventional means such as discing, dragging or mixing operations.

-8-

EXAMPLES

The following Examples are intended to further illustrate the present invention and are not intended to limit the scope of this invention in any manner whatsoever.

Example 1

NMSC was applied (at the rates listed in Table I below) preemergence to aluminum flats (measuring 16 x 23 x 7 cm deep) containing loam soil in which the following species had been sown: *Setaria faberi* (giant foxtail) ("SETFA"); *Sorghum bicolor* (shattercane) ("SORVU"); *Abutilon theophrasti* (velvetleaf) ("ABUTH"); *Amaranthus retroflexus* (redroot pigweed) ("AMARE"); *Chenopodium album* (lambsquarters) ("CHEAL"); *Helianthus annuus* (sunflower) ("HELAN"); *Polygonum pensylvanicum* (Pennsylvania smartweed) ("POLPY"); *Ipomoea purpurea* (tall morningglory) ("PHBPU"); *Xanthium pensylvanicum* (cocklebur) ("XANPE"); *Cyperus esculentus* (yellow nutsedge) ("CYPES"); and corn, *Zea mays* 'Dekalb 656' ("ZEAMX DK656") and 'Garst 8711' ("ZEAMX GA8711"). The soil was fortified with fertilizer (17-17-17) and treated with a fungicide, Captan 80W, prior to seeding. All of the herbicides were applied preemergence with the carrier volume of 234 L/ha (25 gal/A).

After application of NMSC, the flats were placed in a greenhouse. Injury to corn was rated at 8 and 27 days after treatment ("DAT"). Injury was evaluated as percent control, with percent control being the total injury to the plants due to all factors including: inhibited emergence, stunting, malformation, albinism, chlorosis, and other types of plant injury. The control ratings range from 0 to 100 percent, where 0% represents no effect with growth equal to the untreated control and where 100% represents complete kill.

The results observed (as a mean of 3 replications) are summarized in Table I below.

TABLE I

[illegible]

-10-

EXAMPLE 2

Employing a procedure essentially identical to that described in Example 1 except that a silty clay loam soil was employed, NMSC was applied (at the rates listed in Table II below) preemergence to aluminum flats containing the following species: ABUTH; PHBPU; AMARE; POLPY; CYPES; SETFA; *Echinochloa crus-galli* (watergrass) ("ECHCG"); SORVU; and corn hybrids ZEAMX DK656, ZEAMX GA8711, and Pioneer 3475 ("ZEAMX PI3475"). After application, the flats were placed in a greenhouse and the results evaluated after 31 days.

The results observed (as a mean of 3 replications) are summarized in Table II below.

TABLE II

RATE (G/HA)	<u>ABUTH</u>	<u>PHBPU</u>	<u>AMARE</u>	<u>POIPY</u>	<u>CYPES</u>	<u>SETFA</u>	<u>ECICG</u>	<u>SORVU</u>	<u>ZEAMX DK656</u>	<u>ZEAMX GA8711</u>	<u>ZEAMX PI3475</u>
31	98	0	20	80	65	0	33	0	0	0	0
63	100	0	20	73	92	7	7	7	0	0	0
125	80	17	68	83	65	33	0	18	0	0	0
250	100	22	100	98	98	0	100	35	0	0	0
500	100	65	100	98	98	15	100	65	5	0	0

-12-

Example 3

NMSC was applied (at the rates listed in Table III below) preemergence to aluminum flats (measuring 20 x 10 x 6 cm deep) containing soil comprising 2 parts of clay loam to 1 part sandy loam. Prior to application, the following species had been sown: *Ipomea* spp. (morning glory) ("IPOSS"); XANPE; ABUTH; AMARE; *Polygonum lapathifolium* (smartweed) ("POLLA"); *Sesbania exaltata* (sesbania) ("SEBEX"); *Cassia obtusifolia* (sicklepod) ("CASOB"); SETFA; SORVU; ECHCG; CYPES; and corn hybrid 'DeKalb T-1100' ("ZEAMX DK1100"). The flats were placed in a greenhouse and evaluated 17 days after treatment.

The results observed (as a mean of 2 replications) are summarized in Table III below.

TABLE III

RATE	<u>(G/HA)</u>	<u>IPOSS</u>	<u>XANPE</u>	<u>ABUTH</u>	<u>SEBEX</u>	<u>CASOB</u>	<u>POLLA</u>	<u>AMARE</u>	<u>SORVU</u>	<u>CYPES</u>	<u>SETFA</u>	<u>ECHCG</u>	<u>ZEAMX</u> <u>DK1100</u>
125	68	85	100	100	93	45	95	100	88	90	35	95	0
250	83	95	100	100	100	73	100	100	98	90	58	100	0
500	95	100	100	100	100	85	98	100	98	90	78	100	3
1000	100	100	100	100	100	95	98	100	100	93	90	100	35

Example 4

Twelve days after the seeds were sown in aluminum flats (containing soil comprising 2 parts sandy loam to 1 part peat), NMSC was applied (at the rates listed in Table IV) postemergence to plant species listed in Example 3. The flats were placed in a greenhouse and evaluated 17 days after treatment.

The results observed (as a mean of 2 replications) are summarized in Table IV below.

TABLE IV

RATE (G/HA)	<u>IPOSS</u>	<u>XANPE</u>	<u>ABUTH</u>	<u>SEBEX</u>	<u>CASOB</u>	<u>POLLA</u>	<u>AMARE</u>	<u>SORVU</u>	<u>CYPES</u>	<u>SETFA</u>	<u>ECHCG</u>	ZEAMX DK11100
63	100	93	100	100	68	93	100	88	85	55	92	0
125	100	98	100	100	85	100	100	92	90	78	95	2
250	100	93	100	100	93	95	100	95	93	95	100	5
500	100	100	100	100	98	100	100	100	97	98	100	40

-16-

Example 5

NMSC was applied (at the rates listed in Table V) preemergence to aluminum flats (measuring 20x10x6 cm deep) containing soil comprising 1 part clay loam to 1 part sandy loam. Prior to application, the following species had been sown: IPOSS; SORVU; CASOB; SETFA; AMARE; soybean, *Glycine max*, 'Corsoy' ("GLXMA CORSOY"); corn hybrid Stauffer 7751 ("ZEAMX ST7751").

The flats were placed in a greenhouse and evaluated 20 days after treatment. The results observed (as a mean of 2 replications) are summarized in Table V below.

TABLE V

RATE (G/HA)	IPOSS	SORVU	CASOB	SETFA	AMARE	GLXMA CORSOY	ZEAMX ST7751
31	33	0	15	10	98	45	0
63	35	30	15	13	98	55	0
125	63	48	30	35	100	68	0
250	90	85	65	58	98	83	0
500	99	100	83	80	100	93	0

The above results indicate the unexpected corn selectivity of NMSC.

-17-

What is claimed is:

1. A method of selectively controlling undesirable vegetation in corn comprising applying an herbicidally effective amount of 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione to the locus of such vegetation.
2. A method in accordance with claim 1 wherein the 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione is applied at a rate of between about 1 and about 1000 grams per hectare.
3. A method in accordance with claim 2 wherein the 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione is applied at a rate of between about 20 and about 500 grams per hectare.
4. A method in accordance with claim 1 wherein the 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione is applied preemergence.
5. A method in accordance with claim 1 wherein the 2-(2'-nitro-4'-methylsulfonylbenzoyl)-1,3-cyclohexanedione is applied postemergence.

INTERNATIONAL SEARCH REPORT

Inter nal Application No

PC 1, GB 95/02461

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N41/10

According to International Patent Classification (IPC) or to both national classification and IPC:

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR,A,2 675 340 (RHONE POULENC AGRICULTURE) 23 October 1992 see page 2, line 30 - line 31 see page 32; example 3 see page 34; table XV ---	1-5
Y	EP,A,0 186 118 (STAUFFER CHEMICAL CO) 2 July 1986 see page 2, line 4 - line 5 see page 11, table I, comp.no. 26 see page 15; table II see page 17; table III & US,A,5 006 158 cited in the application --- -/--	1-5

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- * & * document member of the same patent family

Date of the actual completion of the international search

30 January 1996

Date of mailing of the international search report

09.02.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

International Application No.

PC, GB 95/02461

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,91 05470 (HOECHST AG) 2 May 1991 see page 1, line 5 - line 33 see page 2, line 1 - page 3, line 4 see page 6, line 11 - line 26 ---	1-5
Y	WO,A,92 19107 (THIELE GERALD H ; THOMAS DAVID A (US)) 12 November 1992 see page 1, paragraph 2 see page 2, paragraph 3 - page 3, paragraph 4 see table 1 ---	1-5
Y	US,A,4 937 386 (UEDA AKIYOSHI ET AL) 26 June 1990 see column 1, line 5 - line 9 see column 1, line 38 - line 40 see claim 2 ---	1-5
A	EP,A,0 563 817 (HOECHST AG) 6 October 1993 see page 2, line 11 - line 19 see page 10, table 1, Bsp. Nr. 13 ---	1-5
A	WO,A,91 05469 (HOECHST AG) 2 May 1991 see page 1, line 5 - line 12 see page 1, line 27 - page 2, line 22 see page 10, line 31 - page 11, line 2 see page 30, line 18 - line 28 -----	1-5

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PC, GB 95/02461

Patent document cited in search report	Publication date	Patent family member(s)	Publicaouon date
FR-A-2675340	23-10-92	CA-A- 2065523	18-10-92
		CA-C- 2065523	18-10-94
		IT-B- 1254876	11-10-95
		US-A- 5416061	16-05-95

EP-A-0186118	02-07-86	AR-A- 240794	28-02-91
		AU-B- 585917	29-06-89
		AU-B- 5133685	26-06-86
		BG-A- 46454	15-12-89
		BG-A- 46452	15-12-89
		CA-A- 1314558	16-03-93
		IE-B- 58400	08-09-93
		JP-B- 7023347	15-03-95
		JP-A- 61152642	11-07-86
		SU-A- 1715190	23-02-92
		US-A- 5006158	09-04-91
		US-A- 4946981	07-08-90

WO-A-9105470	02-05-91	DE-A- 3934706	25-04-91
		DE-A- 3939094	29-05-91
		DE-A- 3943397	04-07-91
		CN-A- 1051840	05-06-91
		CN-A- 1050970	01-05-91
		WO-A- 9105469	02-05-91
		EP-A- 0496751	05-08-92
		EP-A- 0496757	05-08-92
		JP-T- 5501547	25-03-93
		JP-B- 7074126	09-08-95
		JP-T- 5500951	25-02-93
		US-A- 5447903	05-09-95
		US-A- 5318947	07-06-94

WO-A-9219107	12-11-92	EP-A- 0584227	02-03-94

US-A-4937386	26-06-90	JP-A- 2019355	23-01-90

EP-A-0563817	06-10-93	CA-A- 2093105	01-10-93
		JP-A- 6065135	08-03-94

WO-A-9105469	02-05-91	DE-A- 3934706	25-04-91

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/02461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9105469		DE-A- 3939094	29-05-91
		CN-A- 1051840	05-06-91
		CN-A- 1050970	01-05-91
		WO-A- 9105470	02-05-91
		EP-A- 0496751	05-08-92
		EP-A- 0496757	05-08-92
		JP-T- 5501547	25-03-93
		JP-B- 7074126	09-08-95
		JP-T- 5500951	25-02-93
		US-A- 5447903	05-09-95
		US-A- 5318947	07-06-94

THIS PAGE BLANK (USPTO)